

IRSTI: 34.29.25

<https://doi.org/10.70264/jbr.v1.2.2026.3>

**BIOTIC STRESS AND METABOLITES ACCUMULATION IN *MACAIREA RADULA* (BONPL.) DC. (MELASTOMATACEAE) INSECT GALLS**

Patricia Dias Santos\*

Institute of Biology, Uberlândia Federal University, Umuarama Campus, Minas Gerais, Brazil.

\*Corresponding author: Dias Santos P., [diassantos.patricia@gmail.com](mailto:diassantos.patricia@gmail.com)

**ABSTRACT**

Gall formation is known as a new plant organ that depends on biochemical and ecological interactions between the insect and host plant tissues. This relationship is controlled owing biotic stress responses generated by the inducing insect and mediated through the carbohydrates, secondary compounds, and phytohormones biosynthesis in the host plant tissues. The development and growth of galls are associated with primary metabolites, whereas secondary molecules act in defense and ecological functions. These metabolites may change in response to the attack of gall-inducing herbivores, leading to the new compounds biosynthesis and the metals accumulation. This review discusses how host plants react to biotic stress caused by gall-inducing insects.

**Keywords:** Reactive Oxygen Species (ROS); carbohydrates; phenolic compounds.

Received: June 01, 2026 / Accepted: June 26, 2026 / Published: June 27, 2026

© The Author(s) 2026.

**Citation:** Dias Santos P. Biotic stress and metabolites accumulation in *Macairea radula* (Bonpl.) DC. (Melastomataceae) insect galls. Journal of Biological Research, 1(2), 21-26. <https://doi.org/10.70264/jbr.v1.2.2026.3>.

**1. INTRODUCTION**

Plants are affected by a biotic and abiotic stresses variety, which can affect their growth, development, productivity, and geographic distribution [1,2]. To survive adverse environmental conditions, they have developed several adaptive strategies, including the metabolites accumulation that perform protective functions mainly against herbivore attacks [3,4,5]. These organisms induce biotic stress in plants by establishing a species-specific parasitic relationship between hosts and other organisms, such as bacteria, fungi, nematodes, mites, and especially insects [6,7,8,9].

Many of these can alter plant metabolism according to the availability of reactive sites, the plant life cycles synchronicity with the insect during seasonal changes [8,10,11], and even throughout the day [12,13] and induce abnormal growth with cellular hyperplasia and hypertrophy, leading to the formation of newly formed organs, the galls [14,15,16]. These structures can establish themselves in different plant organs such as roots, stems, flower buds, and especially leaves [7,17,15], creating an environment that provides food, shelter, and protection to the gall-inducing insect against natural enemies [18,19].

The stimuli for the induction and development of these organs can be triggered by insect salivary compounds, oviposition, and the phytohormones action [9,20], which are coordinated by gene expression [21,22]. However, the gall structure and metabolism depend on the taxon and, consequently, on

the o gall-inducing insect behavior [8], as well as on how the host tissues and the galled organ respond to oxidative stress [23]. Thus, the structural and metabolic characteristics of galls have been the basis for understanding how the formation and diversification of this new plant organ occur, mainly in the insect group [24,25,26], and how gall-inducing host plants react to biotic stress.

**2. PATTERNS IN GALL DEVELOPMENT**

The galls are like nature ornaments with different shapes, sizes and colors, and modifying the diversity and interactions between organisms. [6]. This diversity has been much studied, mainly regarding the growth processes, morphogenesis, and differentiation [27]. The gall morphotype is “a characteristic neoformed organ generated by the species-specific interaction between an inducing organism and a host plant” [17]. This organ can present conservative patterns, which are determined in plant meristems and manipulated by galling insects.

In the studies of Rohfritsch (1992) [18], gall development has four phases: initiation or induction, growth and development, maturation, and dehiscence or senescence. Some insects are selected to induce galls in young tissues, and this process begins with an increase in cell division and enlargement. In the early phase, there is a predominance of cell hyperplasia, whereas in the later phase cell hypertrophy predominates. During the maturation phase, the insect reaches the last larval instar, which coincides with the highest food consump-

tion. The last phase is senescence or dehiscence, when the insect is ready to emerge and fly. Thus, to explain the adaptive significance of this neof ormation in host plants, there are three main hypotheses: the nutritional, enemy, and the microenvironment hypothesis [19]. The nutritional hypothesis argues that gall tissues offer resources of higher quality and quantity more efficiently to gall-inducing insects than to their free-living ancestors [28], while the microenvironment states that galls protect insects from unfavorable abiotic conditions, such as excessive ultraviolet radiation and temperature, which are major constraints to larval development [19]. According to the enemy hypothesis, the gall protects gall-inducing insects from predators and parasitoids [18, 19].

Neotropical galls are mostly induced by insects of the Cecidomyiidae family, but also by Cynipidae, Psyllidae, Eriophyidae, Apionidae, Gelechioidea, among others [29]. These insects have mouthparts associated with their specific feeding habits, which primarily fall into the chewing, scraping, and sucking guilds [8]. These different feeding activities can trigger differential growth in host tissues [8, 30-32], directly related to the biotic stress establishment [23]. This stress is qualified and quantified through the reactive oxygen species (ROS) presence, generated in plant tissues in response to gall induction [32-34].

Reactive oxygen species result from the molecular oxygen ( $O_2$ ) reduction, forming substances that do not require energy input for reaction, such as superoxide ( $O_2^-$ ), hydrogen peroxide ( $H_2O_2$ ), hydroxyl radical ( $OH\cdot$ ), and singlet oxygen ( $^1O_2$ ) [35]. These molecules trigger oxidative stress in plants and can damage the cell wall, membrane lipids, and even genetic material [36, 37]. However, they can also act as second messengers in many cellular processes, such as transdifferentiation and cellular hypertrophy during the growth and development of newly formed organs [23, 38, 39].

### 3. VASCULARIZATION IN GALLS

The land plants radiation was enabled by the vascular system emergence, which facilitated an increase in plant size and the colonization of diverse ecological niches [40]. This conducting system specialized in the water, mineral solutes, and macromolecules transport differentiates from the procambial meristem into two main tissues: xylem and phloem. The xylem consists of tracheary elements, tracheids and vessel elements characterized as dead, lignified cells at maturity interconnected by perforations that optimize the flow of water and solutes. In contrast, phloem is composed of sieve elements, parenchyma cells, fibers, and sclereids, which are specialized in the translocation of photoassimilates synthesized during photosynthesis.

New organs such as galls are formed in distinct parts of the plant. During this process, photoassimilates are reallocated, and gall initiation and development involve the activity of auxins and cytokinins [41]. Cytokinin is a controlling factor in the regeneration of vessels and sieve tubes around wounds and can influence cell differentiation in vascular tissues. This effect may vary according to cytokinin and auxin physiological levels. In addition, cytokinin may increase vascular cambium sensitivity to auxin stimulation, resulting in a higher phloem/xylem ratio, although these concentrations can vary during gall development. Studies by Mapes and Da-


vies (2001) [42] demonstrated this phenomenon in galls induced by *Eurosta solidaginis* (Diptera) on *Solidago altissima*. Cytokinin was histolocalized in the outer tissue compartment of galls induced by Cecidomyiidae on *Piptadenia gonoacantha* in studies conducted by Bedetti and collaborators (2014) [43]. This finding is one of the strategies discussed regarding how galling insects benefit their life cycle. In addition to this, cytokinin also stimulates the biosynthesis of phenolic compounds, which are partially generated in response to reactive oxygen species (ROS) [23]. Therefore, the local accumulation of phenols, auxins, and ROS may occur simultaneously, resulting in the different shapes and structures of galls. This co-occurrence of molecules can be verified using immunohistochemistry techniques [44].

### 4. HOW THE GALL STRUCTURE WORKS: LOOKING FOR METABOLIC PATHWAYS

In gall development tissues, there are distinct patterns of cell growth and elongation [27]. These patterns can lead to the formation of an external storage compartment and an internal one, the nutritive tissue, directly responsible for the nutrition of the gall-inducing [42-44]. In general, the cells of the nutritive tissue are smaller and less elongated, with dense cytoplasm compared to the cells of the storage tissue, which are composed of hypertrophied and vacuolated cells [32, 42, 44, 45, 47]. These differences are likely associated with the cell wall during gall development [27].

Changes in the cell wall galls composition have been studied to understand microfibril rearrangement and the degree of pectin methylesterification in different host plant systems and gall-inducing insects [27, 41, 48, 49]. Furthermore, these polysaccharides can undergo hydrolysis, releasing oligo- and monosaccharides such as rhamnose (Rha), fucose (Fuc), arabinose (Ara), xylose (Xyl), mannose (Man), glucose (Glc), and galactose (Gal). These carbohydrates are essential components of cell walls and can regulate photosynthetic activity in galls, as well as induce the phenolic compounds biosynthesis, leading to the reddish phenomenon observed in these structures [50,51]. Red galls in *Macairea radula* can accumulate more water-soluble sugars than green galls [52]. To achieve this, the gall-inducing insect continuously stimulates the metabolism of host plants to produce primary metabolites, such as proteins, lipids, and carbohydrates, available not only for the insect's diet [44], but also for the phenolic compounds and polyamines accumulation, which are regulated by phytohormones [53], Figure 1.

All these metabolic changes that begin in the cell wall are directly related to the mechanisms controlling oxidative stress in the gall host plant system. In the galls of *Aspidosperma australe* and *A. spruceanum*, for example, the ROS concentration is found from the initial gall cells until senescence [32]. In fact, the success and establishment of the gall depend on the control of free radical production and elimination through different pathways, both enzymatic and non-enzymatic [23, 53]. One way to dissipate oxidative stress in plants is through the biosynthesis of phenolic compounds, which can donate a hydrogen atom to neutralize ROS activity and prevent cell damage [54-57]. Furthermore, phenolic compounds are capable of sequestering metals, as well as activating antioxidant enzymes such as superoxide dismutase, catalase, ascorbate per-



**Figure 1** –Some metabolites found in galls induced on *M. radula* (Melastomataceae) buds" (Adapted from Santos, 2024 PhD thesis) [45].

oxidases, and peroxidases [58], which act synergistically with phytohormones induced by gall-inducing herbivores and possibly affect the neighboring plant species survival [59, 60].

The cellular machinery used by gall-inducing insects involves not only enzymatic and non-enzymatic systems, but mainly the mediation and regulation by phytohormones [53]. These regulators play a crucial role in the induction, development, and establishment of insect galls, acting as primary chemical messengers [61]. These messengers can stimulate both the primary metabolites production, responsible for maintaining cellular machinery and the nutrition of gall-inducing insects, and secondary compounds, which play a relevant role in maintaining homeostasis, redox reactions, plant defense, and interactions with other organisms [23, 59, 62].

Thus, the entire the gall developmen and its interactions with other organisms can be governed by the action of phytohormones, mainly auxins and cytokinins. The auxin (IAA – indole-3-acetic acid) action has been linked to cell elongation, while cytokinin has been associated with cell division [63]. Other phytohormones, such as abscisic acid, gibberellins, ethylene precursors (ACC), salicylic acid, and methyl jasmonate, as well as polyamines, although participating in responses to abiotic and biotic stresses, are still poorly understood in terms of their functions in the interaction between gall-inducing insects and host plants [53]. It is known that these molecules are associated with the accumulation of metals and phenolic compounds [64, 65].

## 5. FINAL REMARKS

The globoid gall model system induced by *Palaeomystella oligophaga* Becker & Adamski (Lepidoptera) in *Macairea radula* (Bonpl.) DC. (Melastomataceae) has been widely investigated to elucidate the metabolic associations of the insect–plant parasitic interaction. The galls in this system are initiated in the axillary buds and develop as a branched structure on the stem with many leaf-like projections, emergences,

and trichomes [66]. These organs exhibit a typical nutritive tissue that accumulates lipids and proteins [67], and a storage tissue with polysaccharides and phenolic compounds accumulation [52]. The cell walls of the storage and nutritive tissues show differential distribution of pectins and hemicellulose epitopes, evidencing distinct tissue functionalities [53]. Thus, the gall-inducing insect, throughout its different life stages, modifies the gall organ, which assumes different colors ranging from green to red [51,53]. Part of this coloration can be explained by the primary and secondary metabolites [3]. It is believed that various monosaccharides are released within the gall for the biosynthesis of other molecules to maintain antioxidant functions and contribute to the different shapes and colors observed in galls on host plants.

## AUTHORS' CONTRIBUTIONS

**P.S.:** writing – original draft and figure design.

## ACKNOWLEDGEMENT

The author gratefully the FAPEMIG (Fundação de Amparo à Pesquisa do Estado de Minas Gerais) for financial support and fellowship. I am grateful to the Uberlândia Federal University for providing the space and resources necessary for this study.

## FUNDING SOURCES

This research received a doctoral scholarship from FAPEMIG (Fundação de Amparo à Pesquisa do Estado de Minas Gerais).

## CONFLICTS OF INTEREST

The author declares that there are no conflicts of interest.

## OPEN ACCESS

This article is licensed under a Creative Commons Attribution

tion-NonCommercial 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. To view a copy of this licence, visit <https://creativecommons.org/licenses/by-nc/4.0/>.

## REFERENCES

1. Eckardt, N.A., Cutler, S., Juenger, T.E., Marshall-Colon, A., Udvardi, P.E., Verslues, P.E. Focus on climate change and plant abiotic stress biology // *The Plant Cell*. – 2023. – Vol. 35. – P. 1–3. <https://doi.org/10.1093/plcell/koac329>
2. McNichol, B., Russo, S.E. Plant Species' Capacity for Range Shifts at the Habitat and Geographic Scales: A Trade-Off-Based Framework // *Plants*. – 2023. – Vol. 12(6). – P. 1248. <https://doi.org/10.3390/plants12061248>
3. Coelho Kuster, V., Costa Rezende, U., Fernandes Cardoso, J.C., dos Santos Isaias, R.M., Coelho de Oliveira, D. How Galling Organisms Manipulate the Secondary Metabolites in the Host Plant Tissues? A Histochemical Overview in Neotropical Gall Systems. In: Méridon, J.M., Ramawat, K. (eds) Co-Evolution of Secondary Metabolites. Reference Series in Phytochemistry. Springer, Cham. – 2020. – P. 823–842. [https://doi.org/10.1007/978-3-319-96397-6\\_29](https://doi.org/10.1007/978-3-319-96397-6_29)
4. Nicolas-Espinosa, J., Garcia-Ibañez, P., Lopez-Zaplana, A., Yepes-Molina, L., Albaladejo-Marico, L., & Carvajal, M. Confronting Secondary Metabolites with Water Uptake and Transport in Plants under Abiotic Stress // *International Journal of Molecular Sciences*. – 2023. – Vol. 24(3). – P. 2826. <https://doi.org/10.3390/ijms24032826>
5. Ramakrishna, A., Ravishankar, G.A. Influence of abiotic stress signals on secondary metabolites in plants // *Plant Signaling & Behavior*. – 2011. – Vol. 6(11). – P. 1720–1731. <https://doi.org/10.4161/psb.6.11.17613>
6. Isaias, R.M., Carneiro, R.G., Oliveira, D.C., Santos, J.C. Illustrated checklist of Brazilian gall morphotypes // *Neotropical Entomology*. – 2013. – Vol. 42(3). – P. 230–239. <https://doi.org/10.1007/s13744-013-0115-7>
7. Mani, M.S. Introduction to Cecidology, In: J.D. Shorthouse and O. Rohfritsch, Eds. *Biology of Insect-Induced Galls*, Oxford University Press, New York, 1992, pp. 3-7.
8. Oliveira, D.C., Isaias, R.M.S., Fernandes, G.W., Ferreira, B.G., Carneiro, R.G., Fuzaro, L. Manipulation of host plant cells and tissues by gall-inducing insects and adaptive strategies used by different feeding guilds // *Journal of Insect Physiology*. – 2016. – Vol. 84. – P. 103–113. <https://doi.org/10.1016/j.jinsphys.2015.11.012>
9. Shorthouse, J.D., Rohfritsch, O. *Biology of insect-induced galls* // Oxford University Press. – 1992. – P. 285.
10. Pfeffer, L., Rezende, U., Baronio, G.J., Oliveira, D. Building two houses on a single host plant: galling insect synchronizes its life cycle with plant phenology // *Oecologia Australis*. – 2018. – Vol. 22. – P. 438–448. <https://doi.org/10.4257/oeco.2018.2204.07>
11. Yukawa, J. Synchronization of gallers with host plant phenology // *Population Ecology*. – 2000. – Vol. 42. – P. 105–113. <https://doi.org/10.1007/PL00011989>
12. Hirano, T., Okamoto, A., Oda, Y., Sakamoto, T., Takeda, S., Matsuura, T., Ikeda, Y., Higaki, T., Kimura, S., Sato, M. Ab-GALFA: a bioassay for insect gall formation // *Scientific Reports*. – 2023. – Vol. 13(1). – P. 2554. <https://doi.org/10.1038/s41598-023-29302-8>
13. Venkat, A., Muneer, S. Role of circadian rhythms in major plant metabolic and signaling pathways // *Frontiers in Plant Science*. – 2022. – Vol. 13. – P. 836244. <https://doi.org/10.3389/fpls.2022.836244>
14. Harris, M.O., Pitzschke, A. Plants make galls to accommodate foreigners // *New Phytologist*. – 2020. – Vol. 225. – P. 1852–1872. <https://doi.org/10.1111/nph.16340>
15. Mani, M.S. *Ecology of Plant Galls*, Dr. Junk Publisher, The Hague, 1964.
16. Shorthouse, J.D., Wool, D., Raman, A. Gall-inducing insects: nature's most sophisticated herbivores // *Basic and Applied Ecology*. – 2005. – Vol. 6. – P. 407–411. <https://doi.org/10.1016/j.baae.2005.07.001>
17. Isaias, R.M.S., Oliveira, D.C., Carneiro, R., Santos, J. Developmental anatomy of galls in the Neotropics // *Neotropical Insect Galls*. – 2014. – P. 15–34. [https://doi.org/10.1007/978-94-017-8783-3\\_2](https://doi.org/10.1007/978-94-017-8783-3_2)
18. Rohfritsch, O. Patterns in gall development, In: J.D. Shorthouse, O. Rohfritsch, Eds. *Biology of insect induced galls*. Oxford University. Oxford, 1992.
19. Price, P.W., Fernandes, G.W., Waring, G.L. Adaptive nature of insect galls // *Environmental Entomology*. – 1987. – Vol. 16(1). – P. 15–24. <https://doi.org/10.1093/ee/16.1.15>
20. Stone, G.N., Schönrogge, K. The adaptive significance of insect gall morphology // *Trends in Ecology & Evolution*. – 2023. – Vol. 18. – P. 512–522. [https://doi.org/10.1016/S0169-5347\(03\)00247-7](https://doi.org/10.1016/S0169-5347(03)00247-7)
21. Raman, A. Morphogenesis of insect-induced plant galls: fact and questions // *Flora*. – 2011. – Vol. 206(6). – P. 517–533. <https://doi.org/10.1016/j.flora.2010.08.004>
22. Schultz, J.C. et al. A galling insect activates plant reproductive programs during gall development // *Scientific Reports*. – 2019. – Vol. 9(1). – P. 1833. <https://doi.org/10.1038/s41598-018-38475-6>
23. Schultz, J.C., Stone, G.N. A tale of two tissues: probing gene expression in a complex insect-induced gall // *Molecular Ecology*. – 2022. – Vol. 31(11). – P. 3031–3034. <https://doi.org/10.1111/mec.16482>
24. Isaias, R.M.S., Oliveira, D.C., Moreira, A.S.F.P., Soares, G.L.G., Carneiro, R.G. Redox homeostasis in plant galls // *Biochimica et Biophysica Acta*. – 2015. – Vol. 1850. – P. 1509–1517. <https://doi.org/10.1016/j.bbagen.2015.03.007>
25. Julião, G.R., Venticinque, E.M., Fernandes, G.W., Price, P.W. High diversity of galling insects in Amazon // *PLoS ONE*. – 2014. – Vol. 9(12). – P. e114986. <https://doi.org/10.1371/journal.pone.0114986>
26. Maia, V.C., Silva, B.G. Checklist of the gall midges (Diptera, Cecidomyiidae) in the state of Bahia (Northeastern Brazil) / *Lista dos Cecidomyiidae (Diptera) no estado da Bahia (Nordeste do Brasil)* // *Brazilian Journal of Animal and Environmental Research*. – 2020. – Vol. 3(4). – P. 3991–4013. <https://doi.org/10.34188/bjaerv3n4-096>

27. Silva, A.F.M., Oliveira, D.C., Isaias, R.M.S. Influence of the sex of galling insects on gall metabolism: The case of eriococcid galls // *Biochemical Systematics and Ecology*. – 2023. – Vol. 108. – P. 104643. <https://doi.org/10.1016/j.bse.2023.104643>
28. Magalhães, T.A., de Oliveira, D.C., Suzuki, A.Y., Isaias, R.M. Cell elongation patterns in galls // *Protoplasma*. – 2014. – Vol. 251. – P. 747–753. <https://doi.org/10.1007/s00709-013-0574-z>
29. Bronner, R. The role of nutritive cells in the nutrition of cynipids and cecidomyiids // *Biology of Insect-Induced Galls*. – Oxford: Oxford University Press. – 1992. – P. 118–140.
30. Raman, A. Gall induction by hemipteroid insects // *Journal of Plant Interactions*. – 2012. – Vol. 7. – P. 29–44. <https://doi.org/10.1080/17429145.2011.630847>
31. Isaias, R.M.S., Oliveira, D.C., Carneiro, R.G.S. Role of *Euphalerus ostreoides* in manipulating leaflet ontogenesis // *Botany*. – 2011. – Vol. 89. – P. 581–592. <https://doi.org/10.1139/b11-048>
32. Jorge, N.C., Souza-Silva, É.A., Alvarenga, D.R., Saboia, G., Soares, G.L.G., Zini, C.A., Cavalleri, A., Isaias, R.M.S. Structural and chemical profiles of *Myrcia splendens* leaves under gall influence // *Frontiers in Plant Science*. – 2018. – Vol. 9. – P. 1521. <https://doi.org/10.3389/fpls.2018.01521>
33. Oliveira, D.C., Isaias, R.M.S. Redifferentiation of leaflet tissues during gall midrib gall development in *Copaifera langsdorffii* (Fabaceae) // *South African Journal of Botany*. – 2010. – Vol. 76. – P. 239–248. <https://doi.org/10.1016/j.sajb.2009.10.011>
34. Isaias, R.M.S., Oliveira, D. Gall phenotypes product of plant cells defensive responses. In: Mérrillon, J., Ramawat, K. (eds) *Plant Defence: Biological Control. Progress in Biological Control*, vol 12. Springer, Dordrecht. [https://doi.org/10.1007/978-94-007-1933-0\\_11](https://doi.org/10.1007/978-94-007-1933-0_11)
35. Oliveira, D.C., Magalhães, T.A., Carneiro, R.G., Alvim, M.N., Isaias, R.M. Do Cecidomyiidae galls of *Aspidosperma spruceanum* show pre-established cytological and histochemical patterns? // *Protoplasma*. – 2010. – Vol. 242. – P. 81–93. <https://doi.org/10.1007/s00709-010-0128-6>
36. Resende M.L.V., Salgado, S.M.L., Chaves Z.M. Reactive oxygen species in plant defense responses // *Fitopatologia Brasileira*. – 2003. – Vol. 28. – P. 123–130. <https://doi.org/10.1590/S0100-41582003000200001>
37. Demidchik, V. Mechanisms of oxidative stress in plants: from classical chemistry to cell biology // *Environmental and Experimental Botany*. – 2015. – Vol. 109. – P. 212–228. <https://doi.org/10.1016/j.envexpbot.2014.06.021>
38. Meriga, B., Reddy, B.K., Rao, K.R., Reddy, L.A., Kishor P.B.K. Aluminium-induced production of oxygen radicals, lipid peroxidation and DNA damage in seedlings of rice (*Oryza sativa*) // *Journal of Plant Physiology*. – 2004. – Vol 161. – P. 63–68. <https://doi.org/10.1078/0176-1617-01156>
39. Sugimoto, K., Gordon, S.P., Meyerowitz, E.M. Regeneration in plants and animals: dedifferentiation, transdifferentiation, or just differentiation? // *Trends in Cell Biology*. – 2011. – Vol. 21(4). – P. 212–218. <https://doi.org/10.1016/j.tcb.2010.12.004>
40. Yan, J., Tsuchihara, N., Etoh, T., Iwai, S. Reactive oxygen species and nitric oxide in ABA inhibition of stomatal opening // *Plant, Cell & Environment*. – 2007. – Vol. 30. – P. 1320–1325. <https://doi.org/10.1111/j.1365-3040.2007.01711.x>
41. Jorge, N.D.C., Freitas, M.D.S.C., Caffaro, R.M., Vale, F.H.A., Lemos-Filho, J.P., Isaias, R.M.D.S. Vascular traits of stem galls: Cell increment versus morphogenetic constraints in wood anatomy // *Plant Biology*. – 2022. – Vol. 24(3). – P. 450–457. <https://doi.org/10.1111/plb.13392>
42. Mapes, C.C, Davies, P.J. Indole-3-acetic acid and ball gall development on *Solidago altissima*. *New Phytol.* – 2001. – Vol. 151(1). – P. 195–202. <https://doi.org/10.1046/j.1469-8137.2001.00161.x>
43. Bedetti, C.S., Modolo, L.V., dos Santos Isaias, R.M. The role of phenolics in the control of auxin in galls of *Piptadenia gonoacantha* (Mart.) MacBr (Fabaceae: Mimosoideae). *Biochemical Systematics and Ecology*. – 2014. –Vol. 55. – P. 53–59. <https://doi.org/10.1016/j.bse.2014.02.016>
44. Bedetti, C.S., Bragança, G.P., Isaias, R.M.D.S. Influence of auxin and phenolic accumulation on the patterns of cell differentiation in distinct gall morphotypes on *Piptadenia gonoacantha* (Fabaceae). *Australian Journal of Botany*. – 2017. – Vol. 65. – P. 411–420. <https://doi.org/10.1071/BT16257>
45. Santos, Patrícia D. Relation between stress, cell wall alterations, metabolites and metal accumulation in galls induced in *Macairea radula* (Melastomataceae). 172 f. Thesis (PhD in Ecology, Conservation and Biodiversity) – Universidade Federal de Uberlândia, Uberlândia, 2024. <http://doi.org/10.14393/ufu.te.2024.5028>
46. de Oliveira, D.C., Magalhães, T.A., Ferreira, B.G., Teixeira, C.T., Formiga, A.T., Fernandes, G.W., Isaias, R.M. Variation in the degree of pectin methylesterification during the development of *Baccharis dracunculifolia* kidney-shaped gall // *PLoS ONE*. – 2014. – Vol. 9(4). – P. e94588. <https://doi.org/10.1371/journal.pone.0094588>
47. Bragança, G.P.P., Alencar, C.F., Freitas, M.S.C., Isaias, R.M.S. Hemicelluloses and associated compounds determine gall functional traits // *Plant Biology*. – 2020. – Vol. 22. – P. 981–991. <https://doi.org/10.1111/plb.13151>
48. Bragança, O.D.C., Isaias, R.M.S. Compartmentalization of metabolites and enzymatic mediation in nutritive cells of Cecidomyiidae galls on *Piper arboreum* Aubl. // *Journal of Plant Studies*. – 2017. – Vol. 6(11). – P. 11–22. <https://doi.org/10.5539/jps.v6n1p11>
49. Ferreira, B., Freitas, M., Bragança, G., Moreira, A., Carneiro, R., Isaias, R. Enzyme-mediated metabolism in nutritive tissues of galls induced by *Ditylenchus gallaeformans* // *Plant Biology*. – 2019. – Vol. 21(6). – P. 1052–1062. <https://doi.org/10.1111/plb.13009>
50. Ferreira, B.G., Alvarez, R., Avritzer, S.C., Isaias, R.M.S. Revisiting histological patterns of storage tissues // *Botany*. – 2017. – Vol. 95. – P. 173–184. <https://doi.org/10.1139/cjb-2016-0189>
51. Ferreira, B.G., Bragança, G.P., Isaias, R.M.S. Cytological attributes of storage tissues in nematode and eriophyid galls // *Protoplasma*. – 2020. – Vol. 257. – P. 229–244. <https://doi.org/10.1007/s00709-019-01521-1>

doi.org/10.1007/s00709-019-01431-w

52. Isaias, R.M.S., Ferreira, B.G., Alvarenga, D.R., Barbosa, L.R., Salminen, J.P., Steinbauer, J.M. Functional compartmentalisation of nutrients and phenolics in galls // *Austral Entomology*. – 2018. – Vol. 57. – P. 238–246. <https://doi.org/10.1111/aen.12336>

53. Martini, V.C., Moreira, A.S.F.P., Kuster, V.C., Oliveira, D.C. Gallings insects as phenotype manipulators of cell wall composition during the development of galls induced on leaves of *Aspidosperma tomentosum* (Apocynaceae) // *South African Journal of Botany*. – 2019. – Vol. 127. – P. 226–233. <https://doi.org/10.1016/j.sajb.2019.09.006>

54. de Melo Silva, A.F., Lana, L.G., Kuster, V.C., de Oliveira, D.C. Chemical composition of cell wall changes during developmental stages of galls on *Matayba guianensis* (Sapindaceae): perspectives obtained by immunocytochemistry analysis // *Die Naturwissenschaften*. – 2021. – Vol. 108. – P. 108(3), 16. <https://doi.org/10.1007/s00114-021-01732-2>

55. Connor, E.F., Bartlett, L., O'Toole, S., Byrd, S., Biskar, K., Orozco, J. The mechanism of gall induction makes galls red // *Arthropod-Plant Interactions*. – 2012. – Vol. 6. – P. 489–495.

56. Cardoso, J.C.F., Gonçalves, P.H.P., Oliveira, D.C., Rezende, U.C. Host plant intraspecific variation determines gall traits // *Plant Biology*. – 2022. – Vol. 25(1). – P. 208–214. <https://doi.org/10.1111/plb.13472>

57. Naves, G.D.F.S., Kuster, V.C., Machado, M., Santos, P.D., Martini, V.C., Oliveira, D.C. Metabolite investments and stress levels among tissue compartments of *Palaeomystella oligophaga* (Lepidoptera) galls on *Macairea radula* (Melastomataceae) // *Australian Journal of Botany*. – 2021. – Vol. 69. – P. 131–142. <https://doi.org/10.1071/BT20128>

58. Santos, P.D., Bragança, G.P.P., Kuster, V.C., Rocha, D.I., Oliveira, D.C.D. Cell wall epitope distribution in galls induced by *Palaeomystella oligophaga* // *Australian Journal of Botany*. – 2024. – Vol. 72. – P. BT23064. <https://doi.org/10.1071/BT23064>

59. Dumanovic, J., Nepovimova, E., Natic, M., Kuca, K., Jacevic, V. Species and Antioxidant Defense System in Plants: A Concise Overview // *Frontiers in plant science*. – 2021. – Vol. 11. – P. 552969. <https://doi.org/10.3389/fpls.2020.552969>

60. Irato, P., Santovito, G. Enzymatic and non-enzymatic molecules with antioxidant function // *Antioxidants* (Basel, Switzerland). – 2021. – Vol. 10(4). – P. 579. <https://doi.org/10.3390/antiox10040579>

61. Lee, S.H., Ahsan, N., Lee, K.W., Kim, D.H., Lee, D.G., Kwak, S.S., Kwon, S.Y., Kim, T.H., Lee, B.H. Simultaneous overexpression of both CuZn superoxide dismutase and ascorbate peroxidase in transgenic tall fescue plants confers increased tolerance to a wide range of abiotic stresses // *J Plant Physiol*. – 2007. – Vol. 164(12). – P. 1626–1638. <https://doi.org/10.1016/j.jplph.2007.01.003>

62. Sachdev, S. et al. Abiotic stress and reactive oxygen species: generation, signaling, and defense mechanisms // *Antioxidants* (Basel, Switzerland). – 2021. – Vol. 10(2). – P. 277. <https://doi.org/10.3390/antiox10020277>

63. Anjum, N.A., Sharma, P., Gill, S.S., Hasanuzzaman, M., Khan, E.A., Kachhap, K., Mohamed, A.A., Thangavel,

G.D., Devi, P., Vasudhevan, P., Sofo, A., Khan, N.A., Misra, A.N., Lukatkin, A.S., Singh, H.P., Pereira, E., Tuteja, N. Catalase and ascorbate peroxidase-representative H<sub>2</sub>O<sub>2</sub>-detoxifying heme enzymes in plants // *Environmental Science and Pollution Research*. – 2016. – Vol. 23(19). – P. 19002–19029. <https://doi.org/10.1007/s11356-016-7309-6>

64. Baronio, G.J., Oliveira, D.C. Eavesdropping on gall-plant interactions: the importance of the signaling function of induced volatiles // *Plant Signaling & Behavior*. – 2019. – Vol. 14(11). – P. 1665454. <https://doi.org/10.1080/15592324.2019.1665454>

65. Serkedjieva, J. Antioxidant effects of plant polyphenols: A case study on a polyphenol-rich extract from *Geranium sanguineum* L. In: *Reactive Oxygen Species and Antioxidants in Higher Plants*. (eds) Gupta S.D., CRC Press. – 2011. – P. 275–293.

66. Body, M.J.A., Appel, H.M., Edger, P.P., Schultz J.C. WITHDRAWN: A gall-forming insect manipulates hostplant phytohormone synthesis, concentrations, and Signaling // *bioRxiv*. – 2019. – P. 1-72. <https://doi.org/10.1101/658823>

67. Formiga, A.T., Gonçalves, S.J.M.R., Soares, G.L.G., Isaias, R.M.S. Relações entre o teor de fenóis totais e o ciclo das galhas // *Acta Botanica Brasilica*. – 2009. – Vol. 23. – P. 93–99. <https://doi.org/10.1590/S0102-33062009000100012>

68. Sosnowski, J., Truba, M., Vasileva, V. The impact of auxin and cytokinin on growth and development of selected crops // *Agriculture*. – 2023. – Vol. 13. – P. 724. <https://doi.org/10.3390/agriculture13030724>

69. Chen, H., Liu, J., Cui, K., Qin, L., Wang, C., Wu, H., Yang, Z., Ding, W., Shao, S., Wang, H., Ling, X., King-Jones, K., Chen, X. Molecular mechanisms of tannin accumulation in *Rhus* galls and genes involved in plant-insect interactions // *Scientific Reports*. – 2018. – Vol. 8(1). – P. 9841. <https://doi.org/10.1038/s41598-018-28153-y>

70. Tyagi, A., Ali, S., Ramakrishna, G., Singh, A., Park, S., Mahmoudi, H., Bae, H. Revisiting the role of polyamines in plant growth and abiotic stress resilience // *Journal of Plant Growth Regulation*. – 2023. – Vol. 42. – P. 5074–5098. <https://doi.org/10.1007/s00344-022-10847-3>

71. Rezende, U.C., Gonçalves, P.H.P., Kuster, V.C., da Silva Carneiro, R.G., de Oliveira, D.C. Histological features of galls induced on *Macairea radula* allow post-senescence colonization // *CCRN*. – 2023. – Vol. 306. – P. 1-15. <http://dx.doi.org/10.2139/ssrn.4500283>

72. Rezende, U.C., Cardoso, J.C.F., Kuster, V.C., Gonçalves, L.A., Oliveira, D.C. How the activity of natural enemies changes the structure and metabolism of the nutritive tissue in galls? Evidence from the *Palaeomystella oligophaga* (Lepidoptera) -*Macairea radula* (Metastomataceae) system // *Protoplasma*. – 2019. – Vol. 256(3). – P. 669–677. <https://doi.org/10.1007/s00709-018-01339-x>